


a.) Amendment to the Claims:

1. (Currently Amended) ~~An antitussive which comprises, as an active ingredient, a~~ A method for alleviating a cough, which comprises administering, to a patient in need thereof, an effective amount of a pharmaceutical composition comprising a tricyclic compound represented by Formula (I)



{ wherein R¹ represents a hydrogen atom, substituted or unsubstituted lower alkyl, substituted or unsubstituted lower alkoxy or halogen,

X¹-X²-X³ represents ~~CR⁵=CR⁶-CR⁷=CR⁸ [wherein R⁵, R⁶, R⁷ and R⁸ may be the same or different and each represents a hydrogen atom, substituted or unsubstituted lower alkyl, hydroxy, substituted or unsubstituted lower alkoxy, nitro, amino, mono(lower alkyl) substituted amino, di(lower alkyl) substituted amino, substituted or unsubstituted lower alkanoylamino or halogen], N(O)_m=CR⁶-CR⁷=CR⁸ (wherein R⁶, R⁷ and R⁸ have the same meanings as defined above, respectively and m represents 0 or 1), CR⁵=CR⁶-N(O)_m=CR⁸ (wherein R⁵, R⁶, R⁸ and m have the same meanings as defined above, respectively), CR⁵=CR⁶-CR⁷=N(O)_m (wherein R⁵, R⁶, R⁷ and m have the same meanings as defined above, respectively), CR⁵=CR⁶-O (wherein R⁵ and R⁶ have the same meanings~~

~~as defined above, respectively independently represent a hydrogen atom, substituted or unsubstituted lower alkyl, hydroxy, substituted or unsubstituted lower alkoxy, nitro, amino, mono(lower alkyl)-substituted amino, di(lower alkyl)-substituted amino, substituted or unsubstituted lower alkanoylamino or halogen), CR⁵=CR⁶-S (wherein R⁵ and R⁶ have the same meanings as defined above, respectively), O-CR⁷=CR⁸ (wherein R⁷ and R⁸ have the same meanings as R⁵ and R⁶ defined above, respectively), or S-CR⁷=CR⁸ (wherein R⁷ and R⁸ have the same meanings as defined above, respectively) or O-CR⁷=N (wherein R⁷ has the same meaning as defined above),~~


Y represents -CH₂S-, -CH₂SO-, -CH₂SO₂-, -CH₂O-, ~~-CH=CH-~~, (CH₂)_p- (wherein p represents an integer of 0 to 2), -SCH₂-, -SOCH₂-, -SO₂CH₂- or -OCH₂-, and

R² represents a hydrogen atom, substituted or unsubstituted lower alkyl, substituted or unsubstituted lower alkenyl, substituted or unsubstituted lower alkoxy, amino, mono(substituted or unsubstituted lower alkyl)-substituted amino, di(substituted or unsubstituted lower alkyl)-substituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aralkylamino, substituted or unsubstituted arylamino, or a substituted or unsubstituted heterocyclic group}

or a pharmaceutically acceptable salt thereof.

2. (Currently Amended) ~~An antitussive which comprises, as an active ingredient, a~~ A method for alleviating a cough, which comprises administering, to a patient

in need thereof, an effective amount of a pharmaceutical composition comprising a
tricyclic compound represented by Formula (Ia)



[wherein R^1 and X^1 - X^2 - X^3 have the same meanings as defined above,
respectively, represents a hydrogen atom, substituted or unsubstituted lower alkyl,
substituted or unsubstituted lower alkoxy or halogen,

X^1 - X^2 - X^3 represents $CR^5=CR^6-O$ (wherein R^5 and R^6 independently
represent a hydrogen atom, substituted or unsubstituted lower alkyl, hydroxy, substituted
or unsubstituted lower alkoxy, nitro, amino, mono(lower alkyl)-substituted amino, di(lower
alkyl)-substituted amino, substituted or unsubstituted lower alkanoylamino or halogen),

Y^a represents $-CH_2SO_2-$, $-SCH_2-$, $-SOCH_2-$, $-SO_2CH_2-$ ~~or $-OCH_2-$~~ and or
 $-OCH_2-$, and


with the proviso that when Y^a is $-CH_2SO_2-$, $-SCH_2-$, $-SOCH_2-$ or $-SO_2CH_2-$,
then

R^{2a} represents a hydrogen atom, substituted or unsubstituted lower alkyl,
substituted or unsubstituted lower alkenyl, substituted or unsubstituted lower alkoxy,

amino, mono(substituted or unsubstituted lower alkyl)-substituted amino, di(substituted or unsubstituted lower alkyl)-substituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aralkylamino, substituted or unsubstituted arylamino, a substituted or unsubstituted heteroalicyclic group, or a substituted or unsubstituted nitrogen-containing heterocyclic group and

with the proviso that when Y^a is $-OCH_2-$, then

R^{2a} represents a hydrogen atom, trifluoromethyl, substituted or unsubstituted lower alkenyl, substituted or unsubstituted lower alkoxy, amino, mono(substituted or unsubstituted lower alkyl)-substituted amino, di(substituted or unsubstituted lower alkyl)-substituted amino, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted aralkylamino, substituted or unsubstituted arylamino, a substituted or unsubstituted heteroalicyclic group, a substituted or unsubstituted nitrogen-containing heterocyclic group, or Formula (II)



(II)

(wherein n is 0 or 1; R^3 and R^4 may be the same or different and represents a hydrogen atom, substituted or unsubstituted lower alkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted aryl, or substituted or unsubstituted aralkyl, or R^3 and R^4 may be combined together with the adjacent carbon atom thereto to form

cycloalkyl; and Q represents hydroxy, substituted or unsubstituted lower alkoxy, amino or halogen)]

or a pharmaceutically acceptable salt thereof.

3. (Currently Amended) The ~~antitussive~~ method according to Claim 2, wherein Y^a is -CH₂SO₂-, -SCH₂-, -SOCH₂- or -SO₂CH₂-.

4. (Currently Amended) The ~~antitussive~~ method according to Claim 2, wherein Y^a is -OCH₂-.

5. (Currently Amended) The ~~antitussive~~ method according to any of Claims 2 to 4, wherein R¹ is a hydrogen atom, substituted or unsubstituted lower alkoxy or halogen.

6. (Currently Amended) The ~~antitussive~~ method according to any of Claims 2 to 4, wherein R¹ is a hydrogen atom.

7. (Currently Amended) The ~~antitussive~~ method according to claim 2, wherein Y^a is -CH₂SO₂-, -SO₂CH₂- or -OCH₂- and R¹ is a hydrogen atom, substituted or unsubstituted lower alkoxy or halogen.


8. (Currently Amended) The ~~antitussive~~ method according to claim 2, wherein Y^a is -CH₂SO₂- or -SO₂CH₂- and R¹ is a hydrogen atom, substituted or unsubstituted lower alkoxy or halogen.

9. (Currently Amended) The ~~antitussive~~ method according to claim 2, wherein Y^a is -CH₂SO₂- and R¹ is a hydrogen atom, substituted or unsubstituted lower alkoxy or halogen.

10. (Currently Amended) The ~~antitussive~~ method according to any of claims 2 to 4, wherein X¹-X²-X³ is S-CR⁷=CR⁸ (wherein R⁷ and R⁸ ~~have the same meanings as defined above, respectively~~) independently represent a hydrogen atom, substituted or unsubstituted lower alkyl, hydroxy, substituted or unsubstituted lower alkoxy, nitro, amino, mono(lower alkyl)-substituted amino, di(lower alkyl)-substituted amino, substituted or unsubstituted lower alkanoylamino or halogen).

Claim 11 (Cancelled).

12. (Currently Amended) The ~~antitussive~~ method according to any of claims 2 to 4, wherein R^{2a} is Formula (II)



~~(wherein n, R³, R⁴ and Q have the same meanings as defined above, respectively).~~


13. (Currently Amended) The ~~antitussive~~ method according to Claim 12, wherein n is 0.

14. (Currently Amended) The ~~antitussive~~ method according to Claim 13, wherein R³ is methyl, R⁴ is trifluoromethyl, and Q is hydroxy.

15. (Currently Amended) The ~~antitussive~~ method according to Claim 2, wherein R¹ is a hydrogen atom, Y^a is -CH₂SO₂-, X¹-X²-X³ is S-CR⁷=CR⁸ (wherein R⁷ and R⁸ ~~have the same meanings as defined above, respectively~~) independently represent a


hydrogen atom, substituted or unsubstituted lower alkyl, hydroxy, substituted or unsubstituted lower alkoxy, nitro, amino, mono(lower alkyl)-substituted amino, di(lower alkyl)-substituted amino, substituted or unsubstituted lower alkanoylamino or halogen),
and

R² is Formula (III)



(III)

16. (Currently Amended) ~~An antitussive which comprises, as an active ingredient,~~ A method for alleviating of a cough, which comprises administering, to a patient in need thereof, an effective amount of a pharmaceutical composition comprising a
tricyclic compound represented by Formula (Ib)




(Ib)

[wherein R^1 and $X^1-X^2-X^3$ have the same meanings as defined above,
respectively represents a halogen atom, substituted or unsubstituted lower alkyl,
substituted or unsubstituted lower alkoxy or halogen,

$X^1-X^2-X^3$ represents $CR^5=CR^6-O$, $CR^5=CR^6-S$, $O-CR^7=CR^8$, or $S-CR^7+CR^8$
(wherein R^5 and R^6 independently represent a hydrogen atom, substituted or unsubstituted
lower alkyl, hydroxy, substituted or unsubstituted lower alkoxy, nitro, amino, mono(lower
alkyl)-substituted amino, di(lower alkyl)-substituted amino, substituted or unsubstituted
lower alkanoylamino or halogen, and R^7 and R^8 have the same meaning as R^5 and R^6 ,
respectively,

Y^b represents ~~$-CH_2O-$, $-CH_2S-$, $-CH_2SO-$, $CH=CH$ or $-(CH_2)_p-$ (wherein p~~
~~has the same meaning as defined above)~~ $-CH_2O-$, $-CH_2S-$ or $-CH_2SO-$, and

R^{2b} represents Formula (III)



or a pharmaceutically acceptable salt thereof.

Claim 17 (Cancelled).

18. (Currently Amended) The ~~antitussive~~ method according to Claim 16, wherein X^1 - X^2 - X^3 is $CR^5=CR^6-O$ (~~wherein R^5 and R^6 have the same meanings as defined above, respectively~~) or $CR^5=CR^6-S$ (~~wherein R^5 and R^6 have the same meanings as defined above, respectively~~).

19. (Currently Amended) The ~~antitussive~~ method according to Claim 16, wherein X^1 - X^2 - X^3 is $O-CR^7=CR^8$ (~~wherein R^7 and R^8 have the same meanings as defined above, respectively~~) or $S-CR^7=CR^8$ (~~wherein R^7 and R^8 have the same meanings as defined above, respectively~~).

20. (Currently Amended) The ~~antitussive~~ method according to any of ~~Claims 16 to 19~~ Claims 16, 18 and 19, wherein Y^b is $-CH_2O-$.

Claims 21-24 (Cancelled).

25. (Currently Amended) The ~~antitussive~~ method according to any of ~~Claims 16 to 19~~ Claims 16, 18 and 19, wherein Y^b is $-CH_2S-$ or $-CH_2SO-$.

Claims 26-27 (Cancelled).